

# Package ‘ntsDists’

May 9, 2026

**Type** Package

**Title** Neutrosophic Distributions

**Version** 2.1.1

**Maintainer** Danial Mazarei <daniel.mazarei@gmail.com>

**Description** Computes the pdf, cdf, quantile function and generating random numbers for neutrosophic distributions. This family have been developed by different authors in the recent years. See Patro and Smarandache (2016) <[doi:10.5281/zenodo.571153](https://doi.org/10.5281/zenodo.571153)> and Rao et al (2023) <[doi:10.5281/zenodo.7832786](https://doi.org/10.5281/zenodo.7832786)>.

**BugReports** <https://github.com/dmazarei/ntsDists/issues>

**License** GPL (>= 2)

**URL** <https://github.com/dmazarei/ntsDists>

**Encoding** UTF-8

**LazyData** false

**RoxygenNote** 7.2.3

**Depends** R (>= 3.5)

**NeedsCompilation** no

**Author** Danial Mazarei [aut, cre] (ORCID: <<https://orcid.org/0000-0002-3633-9298>>),  
Mina Norouzirad [aut] (ORCID: <<https://orcid.org/0000-0003-0311-6888>>),  
Amin Roshani [aut] (ORCID: <<https://orcid.org/0000-0002-3329-5330>>),  
Gadde Srinivasa Rao [ctb] (ORCID: <<https://orcid.org/0000-0002-3683-5486>>),  
Foad Esmaeili [ctb] (ORCID: <<https://orcid.org/0000-0002-9638-0807>>),  
FCT, I.P. [fnd] (under the scope of the projects UIDB/00297/2020 and UIDP/00297/2020 (NovaMath))

**Repository** CRAN

**Date/Publication** 2024-04-17 07:00:02 UTC

## Contents

balls . . . . .	2
Neutrosophic Beta . . . . .	3
Neutrosophic Binomial . . . . .	4
Neutrosophic Discrete Uniform . . . . .	6
Neutrosophic Exponential . . . . .	7
Neutrosophic Gamma . . . . .	9
Neutrosophic Generalized Exponential . . . . .	10
Neutrosophic Generalized Pareto . . . . .	11
Neutrosophic Generalized Rayleigh . . . . .	13
Neutrosophic Geometric . . . . .	14
Neutrosophic Kumaraswamy . . . . .	16
Neutrosophic Laplace . . . . .	17
Neutrosophic Negative Binomial . . . . .	19
Neutrosophic Normal . . . . .	20
Neutrosophic Poisson . . . . .	21
Neutrosophic Rayleigh . . . . .	23
Neutrosophic Uniform . . . . .	24
Neutrosophic Weibull . . . . .	26
remission . . . . .	27
<b>Index</b>	<b>28</b>

---

balls

*Balls data*

---

### Description

It is related to failure times of 23 bearing balls.

### Format

A data.frame with 23 observations of failure times of bearing balls.

### Source

Lawless, J. F. (2003). *Statistical Models and Methods for Lifetime Data*, Wiley, Hoboken, NJ, USA.  
 Salam, S., Khan, Z., Ayed, H., Brahmia, A., Amin, A. (2021). The Neutrosophic Lognormal Model in Lifetime Data Analysis: Properties and Applications, *Fuzzy Sets and Their Applications in Mathematics*, Article ID 6337759.

### Examples

```
data("balls")
balls
```

**Description**

Density, distribution function, quantile function and random generation for the neutrosophic Beta distribution with shape parameters  $\text{shape1} = \alpha_N$  and  $\text{shape2} = \beta_N$ .

**Usage**

```
dnsBeta(x, shape1, shape2)
pnsBeta(q, shape1, shape2, lower.tail = TRUE)
qnsBeta(p, shape1, shape2)
rnsBeta(n, shape1, shape2)
```

**Arguments**

<code>x</code>	a vector or matrix of observations for which the pdf needs to be computed.
<code>shape1</code>	the first shape parameter, which must be a positive interval.
<code>shape2</code>	the second shape parameter, which must be a positive interval.
<code>q</code>	a vector or matrix of quantiles for which the cdf needs to be computed.
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
<code>p</code>	a vector or matrix of probabilities for which the quantile needs to be computed.
<code>n</code>	number of random values to be generated.

**Details**

The neutrosophic beta distribution with parameters  $\alpha_N$  and  $\beta_N$  has the probability density function

$$f_N(x) = \frac{1}{B(\alpha_N, \beta_N)} x^{\alpha_N-1} (1-x)^{\beta_N-1}$$

for  $\alpha_N \in (\alpha_L, \alpha_U)$ , the first shape parameter which must be a positive interval, and  $\beta_N \in (\beta_L, \beta_U)$ , the second shape parameter which must also be a positive interval, and  $0 \leq x \leq 1$ . The function  $B(a, b)$  returns the beta function and can be calculated using [beta](#).

**Value**

`dnsBeta` gives the density function  
`pnsBeta` gives the distribution function  
`qnsBeta` gives the quantile function  
`rnsBeta` generates random values from the neutrosophic Beta distribution.

## References

Sherwani, R. Ah. K., Naeem, M., Aslam, M., Reza, M. A., Abid, M., Abbas, S. (2021). Neutrosophic beta distribution with properties and applications. *Neutrosophic Sets and Systems*, 41, 209-214.

## Examples

```
dnsBeta(x = c(0.1, 0.2), shape1 = c(1, 1), shape2 = c(2, 2))
dnsBeta(x = 0.1, shape1 = c(1, 1), shape2 = c(2, 2))

x <- matrix(c(0.1, 0.1, 0.2, 0.3, 0.5, 0.5), ncol = 2, byrow = TRUE)
dnsBeta(x, shape1 = c(1, 2), shape2 = c(2, 3))

pnsBeta(q = c(0.1, 0.1), shape1 = c(3, 1), shape2 = c(1, 3), lower.tail = FALSE)
pnsBeta(x, shape1 = c(1, 2), shape2 = c(2, 2))

qnsBeta(p = 0.1, shape1 = c(1, 1), shape2 = c(2, 2))
qnsBeta(p = c(0.25, 0.5, 0.75), shape1 = c(1, 2), shape2 = c(2, 2))

# Simulate 10 numbers
rnsBeta(n = 10, shape1 = c(1, 2), shape2 = c(1, 1))
```

---

Neutrosophic Binomial *Neutrosophic Binomial Distribution*

---

## Description

Density, distribution function, quantile function and random generation for the neutrosophic binomial distribution with parameters size =  $n$  and prob =  $p_N$ .

## Usage

```
dnsBinom(x, size, prob)

pnsBinom(q, size, prob, lower.tail = TRUE)

qnsBinom(p, size, prob)

rnsBinom(n, size, prob)
```

## Arguments

x	a vector or matrix of observations for which the pdf needs to be computed.
size	number of trials (zero or more), which must be a positive interval.
prob	probability of success on each trial, $0 \leq prob \leq 1$ .

q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

### Details

The neutrosophic binomial distribution with parameters  $n$  and  $p_N$  has the density

$$f_X(x) = \binom{n}{x} p_N^x (1 - p_N)^{n-x}$$

for  $n \in \{1, 2, \dots\}$  and  $p_N \in (p_L, p_U)$  which must be  $0 < p_N < 1$  and  $x \in \{0, 1, 2, \dots, n\}$ .

### Value

dnsBinom gives the probability mass function

pnsBinom gives the distribution function

qnsBinom gives the quantile function

rnsBinom generates random variables from the Binomial Distribution.

### References

Granados, C. (2022). Some discrete neutrosophic distributions with neutrosophic parameters based on neutrosophic random variables. *Hacetatepe Journal of Mathematics and Statistics*, 51(5), 1442-1457.

### Examples

```
# Probability of X = 17 when X follows bin(n = 20, p = [0.9,0.8])
dnsBinom(x = 17, size = 20, prob = c(0.9, 0.8))

x <- matrix(c(15, 15, 17, 18, 19, 19), ncol = 2, byrow = TRUE)
dnsBinom(x = x, size = 20, prob = c(0.8, 0.9))

pnsBinom(q = 17, size = 20, prob = c(0.9, 0.8))
pnsBinom(q = c(17, 18), size = 20, prob = c(0.9, 0.8))
pnsBinom(q = x, size = 20, prob = c(0.9, 0.8))

qnsBinom(p = 0.5, size = 20, prob = c(0.8, 0.9))
qnsBinom(p = c(0.25, 0.5, 0.75), size = 20, prob = c(0.8, 0.9))

# Simulate 10 numbers
rnsBinom(n = 10, size = 20, prob = c(0.8, 0.9))
```

---

 Neutrosophic Discrete Uniform

*Neutrosophic Discrete Uniform Distribution*


---

### Description

Density, distribution function, quantile function and random generation for the neutrosophic discrete uniform distribution with parameter  $k_N$ .

### Usage

```
dnsDiscUnif(x, k)
pnsDiscUnif(q, k, lower.tail = TRUE)
qnsDiscUnif(p, k)
rnsDiscUnif(n, k)
```

### Arguments

x	a vector or matrix of observations for which the pdf needs to be computed.
k	parameter of the distribution that must be a positive finite interval.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

### Details

Let  $X_N$  be a neutrosophic random variable and denote  $X_N \sim \mathcal{DU}(1, 2, \dots, k_N)$  as neutrosophic discrete uniform distribution with parameter  $k_N$  has the density

$$f_N(x) = \frac{1}{k_N}$$

for  $k_N \in (k_L, k_U)$ .

### Value

dnsDiscUnif gives the probability mass function,  
 pnsDiscUnif gives the distribution function  
 qnsDiscUnif gives the quantile function  
 rnsDiscUnif generates random variables from the neutrosophic Discrete Uniform Distribution.

## References

Granados, C. (2022). Some discrete neutrosophic distributions with neutrosophic parameters based on neutrosophic random variables. *Hacetatepe Journal of Mathematics and Statistics*, 51(5), 1442-1457.

## Examples

```
dnsDiscUnif(x = 8, k = c(10, 11))
dnsDiscUnif(x = c(8, 9), k = c(10, 11))

pnsDiscUnif(q = 2, k = c(10, 11))

qnsDiscUnif(p = 0.2, k = c(10, 11))

# Simulate 10 numbers
rnsDiscUnif(n = 10, k = c(10, 11))
```

---

Neutrosophic Exponential

*Neutrosophic Exponential Distribution*

---

## Description

Density, distribution function, quantile function and random generation for the neutrosophic exponential distribution with the parameter rate =  $\theta_N$ .

## Usage

```
dnsExp(x, rate)

pnsExp(q, rate, lower.tail = TRUE)

qnsExp(p, rate)

rnsExp(n, rate)
```

## Arguments

x	a vector or matrix of observations for which the pdf needs to be computed.
rate	the shape parameter, which must be a positive interval.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

**Details**

The neutrosophic exponential distribution with parameter  $\theta_N$  has density

$$f_N(x) = \theta_N \exp(-x\theta_N)$$

for  $x \geq 0$  and  $\theta_N \in (\theta_L, \theta_U)$ , the rate parameter must be a positive interval and  $x \geq 0$ .

**Value**

dnsExp gives the density function

pnsExp gives the distribution function

qnsExp gives the quantile function

rnsExp generates random values from the neutrosophic exponential distribution.

**References**

Duan, W., Q., Khan, Z., Gulistan, M., Khurshid, A. (2021). Neutrosophic Exponential Distribution: Modeling and Applications for Complex Data Analysis, *Complexity*, 2021, 1-8.

**Examples**

```
# Example 4 of Duan et al. (2021)

data <- matrix(c(4, 4, 3.5, 3.5, 3.9, 4.1, 4.2, 4.2, 4.3, 4.6, 4.7, 4.7),
  nrow = 6, ncol = 2, byrow = TRUE)

dnsExp(data, rate = c(0.23, 0.24))
dnsExp(x = c(4, 4.1), rate = c(0.23, 0.24))

dnsExp(4, rate = c(0.23, 0.23))

# The cumulative distribution function for the neutrosophic observation (4,4.1)
pnsExp(c(4, 4.1), rate = c(0.23, 0.24), lower.tail = TRUE)

pnsExp(4, rate = c(0.23, 0.24))
# The first percentile
qnsExp(p = 0.1, rate = 0.25)

# The quantiles
qnsExp(p = c(0.25, 0.5, 0.75), rate = c(0.24, 0.25))

# Simulate 10 numbers
rnsExp(n = 10, rate = c(0.23, 0.24))
```

**Description**

Density, distribution function, quantile function and random generation for the neutrosophic gamma distribution with parameter shape =  $\alpha_N$  and scale =  $\lambda_N$ .

**Usage**

```
dnsGamma(x, shape, scale)
pnsGamma(q, shape, scale, lower.tail = TRUE)
qnsGamma(p, shape, scale)
rnsGamma(n, shape, scale)
```

**Arguments**

x	a vector or matrix of observations for which the pdf needs to be computed.
shape	the shape parameter, which must be a positive interval.
scale	the scale parameter, which must be a positive interval.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

**Details**

The neutrosophic gamma distribution with parameters  $\alpha_N$  and  $\lambda_N$  has density

$$f_N(x) = \frac{1}{\Gamma(\alpha_N)\lambda_N^{\alpha_N}} x^{\alpha_N-1} \exp\{- (x/\lambda_N)\}$$

for  $x \geq 0$ ,  $\alpha_N \in (\alpha_L, \alpha_U)$ , the shape parameter which must be a positive interval and  $\lambda_N \in (\lambda_L, \lambda_U)$ , the scale parameter which must be a positive interval. Here,  $\Gamma(\cdot)$  is gamma function implemented by [gamma](#).

**Value**

dnsGamma gives the density function  
 pnsGamma gives the distribution function  
 qnsGamma gives the quantile function  
 rnsGamma generates random variables from the neutrosophic gamma distribution.

## References

Khan, Z., Al-Bossly, A., Almazah, M. M. A., and Alduais, F. S. (2021). On statistical development of neutrosophic gamma distribution with applications to complex data analysis, *Complexity*, 2021, Article ID 3701236.

## Examples

```
data(remission)
dnsGamma(x = remission, shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))

pnsGamma(q = 20, shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))

# Calculate quantiles
qnsGamma(p = c(0.25, 0.5, 0.75), shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))

# Simulate 10 numbers
rnsGamma(n = 10, shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))
```

---

Neutrosophic Generalized Exponential

*Neutrosophic Generalized Exponential Distribution*

---

## Description

Density, distribution function, quantile function and random generation for the neutrosophic generalized exponential distribution with shape parameter  $\delta_N$  and scale parameter  $\nu_N$ .

## Usage

```
dnsGenExp(x, nu, delta)

pnsGenExp(q, nu, delta, lower.tail = TRUE)

qnsGenExp(p, nu, delta)

rnsGenExp(n, nu, delta)
```

## Arguments

x	a vector or matrix of observations for which the pdf needs to be computed.
nu	the scale parameter, which must be a positive interval.
delta	the shape parameter, which must be a positive interval.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

**Details**

The neutrosophic generalized exponential distribution with parameters  $\delta_N$  and  $\nu_N$  has density

$$f_N(x) = \frac{\delta_N}{\nu_N} \left( 1 - \exp \left\{ -\frac{x_N}{\nu_N} \right\} \right)^{\delta_N - 1} \exp \left\{ -\frac{x_N}{\nu_N} \right\}$$

for  $\delta_N \in (\delta_L, \delta_U)$ , the shape parameter which must be a positive interval, and  $\nu_N \in (\nu_L, \nu_U)$ , the scale parameter which must also be a positive interval, and  $x \geq 0$ .

**Value**

dnsGenExp gives the density function

pnsGenExp gives the distribution function

qnsGenExp gives the quantile function

rnsGenExp generates random variables from the neutrosophic generalized exponential distribution.

**References**

Rao, G. S., Norouzirad, M., and Mazarei . D. (2023). Neutrosophic Generalized Exponential Distribution with Application. *Neutrosophic Sets and Systems*, 55, 471-485.

**Examples**

```
data(remission)
dnsGenExp(x = remission, nu = c(7.9506, 8.0568), delta = c(1.2390, 1.2397))

pnsGenExp(q = 20, nu = c(7.9506, 8.0568), delta = c(1.2390, 1.2397))

# Calcluate quantiles
qnsGenExp(c(0.25, 0.5, 0.75), nu = c(7.9506, 8.0568), delta = c(1.2390, 1.2397))

# Simulate 10 values
rnsGenExp(n = 10, nu = c(7.9506, 8.0568), delta = c(1.2390, 1.2397))
```

---

Neutrosophic Generalized Pareto

*Neutrosophic Generalized Pareto Distribution*

---

**Description**

Density, distribution function, quantile function and random generation for the neutrosophic generalized pareto distribution with parameters shape =  $\alpha_N$  and scale =  $\beta_N$ .

**Usage**

```

dnsGenPareto(x, shape, scale)

pnsGenPareto(q, shape, scale, lower.tail = TRUE)

qnsGenPareto(p, shape, scale)

rnsGenPareto(n, shape, scale)

```

**Arguments**

`x` a vector or matrix of observations for which the pdf needs to be computed.

`shape` the shape parameter, which must be a positive interval.

`scale` the scale parameter, which must be a positive interval.

`q` a vector or matrix of quantiles for which the cdf needs to be computed.

`lower.tail` logical; if TRUE (default), probabilities are  $P(X \leq x)$ ; otherwise,  $P(X > x)$ .

`p` a vector or matrix of probabilities for which the quantile needs to be computed.

`n` number of random values to be generated.

**Details**

The neutrosophic generalized pareto distribution with parameters  $\alpha_N$  and  $\beta_N$  has density

$$f_N(x) = \frac{1}{\beta_N} \left(1 + \frac{\alpha_N x_N}{\beta_N}\right)^{-\frac{1}{\alpha_N} - 1}$$

for  $x \geq 0$ ,  $\alpha_N \in (\alpha_L, \alpha_U)$ , the shape parameter which must be a positive interval and  $\beta_N \in (\beta_L, \beta_U)$ , the scale parameter which must be a positive interval.

**Value**

`dnsGenPareto` gives the density function

`pnsGenPareto` gives the distribution function

`qnsGenPareto` gives the quantile function

`rnsGenPareto` generates random variables from the neutrosophic generalized pareto distribution.

**References**

Eassa, N. I., Zaher, H. M., & El-Magd, N. A. A. (2023). Neutrosophic Generalized Pareto Distribution, *Mathematics and Statistics*, 11(5), 827–833.

**Examples**

```

data(remission)
dnsGenPareto(x = remission, shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))

pnsGenPareto(q = 20, shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))

# Calculate quantiles
qnsGenPareto(p = c(0.25, 0.5, 0.75), shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))

# Simulate 10 numbers
rnsGenPareto(n = 10, shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))

```

---

Neutrosophic Generalized Rayleigh

*Neutrosophic Generalized Rayleigh Distribution*


---

**Description**

Density, distribution function, quantile function and random generation for the neutrosophic generalized Rayleigh distribution with parameters shape =  $\nu_N$  and scale =  $\sigma_N$ .

**Usage**

```

dnsGenRayleigh(x, shape, scale)

pnsGenRayleigh(q, shape, scale, lower.tail = TRUE)

qnsGenRayleigh(p, shape, scale)

rnsGenRayleigh(n, shape, scale)

```

**Arguments**

x	a vector or matrix of observations for which the pdf needs to be computed.
shape	the shape parameter, which must be a positive interval.
scale	the scale parameter, which must be a positive interval.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

**Details**

The neutrosophic generalized Rayleigh distribution with parameters  $\nu_N$  and  $\sigma_N$  has the density

$$f_N(x) = \frac{2\nu_N}{\sigma_N^2} x \exp\left\{-\left(\frac{x}{\sigma_N}\right)^2\right\} \left[1 - \exp\left\{-\left(\frac{x}{\sigma_N}\right)^2\right\}\right]^{\nu_N-1}$$

for  $x > 0$ ,  $\nu_N \in (\nu_L, \nu_U)$ , the shape parameter which must be a positive interval and  $\sigma_N \in (\sigma_L, \sigma_U)$ , the scale parameter which must be a positive interval.

**Value**

`dnsGenRayleigh` gives the density function

`pnsGenRayleigh` gives the distribution function

`qnsGenRayleigh` gives the quantile function

`rnsGenRayleigh` generates random variables from the Neutrosophic Generalized Rayleigh Distribution.

**References**

Norouzirad, M., Rao, G. S., & Mazarei, D. (2023). Neutrosophic Generalized Rayleigh Distribution with Application. *Neutrosophic Sets and Systems*, 58(1), 250-262.

**Examples**

```
data(remission)
dnsGenRayleigh(x = remission, shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))

pnsGenRayleigh(q = 20, shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))

# Calculate quantiles
qnsGenRayleigh(p = c(0.25, 0.5, 0.75), shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))

# Simulate 10 values
rnsGenRayleigh(n = 10, shape = c(1.1884, 1.1896), scale = c(7.6658, 7.7796))
```

---

Neutrosophic Geometric

*Neutrosophic Geometric Distribution*

---

**Description**

Density, distribution function, quantile function and random generation for the neutrosophic Geometric distribution with parameter  $\text{prob} = p_N$ .

**Usage**

```

dnsGeom(x, prob)

pnsGeom(q, prob, lower.tail = TRUE)

qnsGeom(p, prob)

rnsGeom(n, prob)

```

**Arguments**

x	a vector or matrix of observations for which the pdf needs to be computed.
prob	probability of success on each trial, $\text{prob} \in (0, 1)$ .
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

**Details**

The neutrosophic Geometric distribution with parameter  $p_N$  has the density

$$f_X(x) = p_N (1 - p_N)^x$$

for  $p_N \in (p_L, p_U)$  which must be  $0 < p_N < 1$  and  $x \in \{0, 1, 2, \dots\}$ .

**Value**

dnsGeom gives the probability mass function  
pnsGeom gives the distribution function  
qnsGeom gives the quantile function  
rnsGeom generates random variables from the Geometric Distribution.

**References**

Granados, C. (2022). Some discrete neutrosophic distributions with neutrosophic parameters based on neutrosophic random variables. *Hacetatepe Journal of Mathematics and Statistics*, 51(5), 1442-1457.

**Examples**

```

# One person participates each week with a ticket in a lottery game, where
# the probability of winning the first prize is (10^(-8), 10^(-6)).
# Probability of one persons wins at the fifth year?

dnsGeom(x = 5, prob = c(1e-8, 1e-6))

# Probability of one persons wins after 10 years?

```

```

pnsGeom(q = 10, prob = c(1e-8, 1e-6))
pnsGeom(q = 10, prob = c(1e-8, 1e-6), lower.tail = FALSE)
# Calculate the quantiles
qnsGeom(p = c(0.25, 0.5, 0.75), prob = c(1e-8, 1e-6))
# Simulate 10 numbers
rnsGeom(n = 10, prob = c(1e-8, 1e-6))

```

---

Neutrosophic Kumaraswamy

*Neutrosophic Kumaraswamy Distribution*

---

### Description

Density, distribution function, quantile function and random generation for the neutrosophic Kumaraswamy distribution with shape parameters  $\alpha_N$  and  $\beta_N$ .

### Usage

```

dnsKumaraswamy(x, shape1, shape2)

pnsKumaraswamy(q, shape1, shape2, lower.tail = TRUE)

qnsKumaraswamy(p, shape1, shape2)

rnsKumaraswamy(n, shape1, shape2)

```

### Arguments

x	a vector or matrix of observations for which the pdf needs to be computed.
shape1	the shape parameter, which must be a positive interval.
shape2	the shape parameter, which must be a positive interval.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

### Details

The neutrosophic Kumaraswamy distribution with parameters  $\alpha_N$  and  $\beta_N$  has density

$$f_N(x) = \alpha_N \beta_N x^{\alpha_N - 1} (1 - x^{\alpha_N})^{\beta_N - 1}$$

for  $0 \leq x \leq 1$ ,  $\alpha_N \in (\alpha_L, \alpha_U)$  and  $\beta_N \in (\beta_L, \beta_U)$  are shape parameters.

**Value**

pnsKumaraswamy gives the distribution function  
 dnsKumaraswamy gives the density  
 qnsKumaraswamy gives the quantile function  
 rnsKumaraswamy generates random values from the neutrosophic Kumaraswamy distribution.

**References**

Ahsan-ul-Haq, M. (2022). Neutrosophic Kumaraswamy Distribution with Engineering Application, *Neutrosophic Sets and Systems*, 49, 269-276.

**Examples**

```
dnsKumaraswamy(x = c(0.5, 0.1), shape1 = c(0.23, 0.24), shape2 = c(1, 2))
dnsKumaraswamy(0.5, shape1 = c(0.23, 0.24), shape2 = c(1, 2))

# The cumulative distribution function for the neutrosophic observation (4,4.1)
pnsKumaraswamy(q = c(.8, .1), shape1 = c(0.23, 0.24), shape2 = c(1, 2))
# The first percentile
qnsKumaraswamy(p = 0.1, shape1 = 0.24, shape2 = 2)

# The quantiles
qnsKumaraswamy(p = c(0.25, 0.5, 0.75), shape1 = c(0.23, 0.24), shape2 = c(1, 2))

# Simulate 10 numbers
rnsKumaraswamy(n = 10, shape1 = c(0.23, 0.24), shape2 = c(1, 2))
```

---

Neutrosophic Laplace    *Neutrosophic Laplace (Double Exponential) Distribution*

---

**Description**

Density, distribution function, quantile function, and random generation for the neutrosophic Laplace (Double Exponential) distribution with parameters location =  $\theta_N$  and scale =  $\beta_N$ .

**Usage**

```
dnsLaplace(x, location, scale)

pnsLaplace(q, location, scale, lower.tail = TRUE)

qnsLaplace(p, location, scale)

rnsLaplace(n, location, scale)
```

**Arguments**

x	a vector or matrix of observations for which the pdf needs to be computed.
location	the location parameter, which is the mean.
scale	the scale parameter, Must consist of positive values.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

**Details**

The neutrosophic Laplace distribution with parameters  $\theta_N$  and  $\beta_N$  has density

$$f_N(x) = \frac{1}{2\beta_N} \exp\left\{-\frac{|x - \theta_N|}{\beta_N}\right\}$$

for  $-\infty < x < \infty$ ,  $\theta_N \in (\theta_L, \theta_U)$ , the location parameter,  $\beta_N \in (\beta_L, \beta_U)$ , the scale parameter which be a positive interval.

**Value**

dnsLaplace gives the density function

pnsLaplace gives the distribution function

qnsLaplace gives the quantile function

rnsLaplace generates random values from the neutrosophic Laplace distribution.

**References**

Rahul, T., Malik, S. C., Raj, M. (2023). Neutrosophic Laplace Distribution with Application in Financial Data Analysis, *Neutrosophic Sets and Systems*, 57(1), 224-233.

**Examples**

```
dnsLaplace(x = c(4, 4.1), location = c(0.23, 0.24), scale = c(1, 2))
dnsLaplace(4, location = c(0.23, 0.24), scale = c(1, 2))
```

```
# The cumulative distribution function for the neutrosophic observation (4,4.1)
pnsLaplace(q = c(4, 4.1), location = c(0.23, 0.24), scale = c(1, 2))
# The first percentile
qnsLaplace(p = 0.1, location = 0.24, scale = 2)
```

```
# The quantiles
qnsLaplace(p = c(0.25, 0.5, 0.75), location = c(0.23, 0.24), scale = c(1, 2))
```

```
# Simulate 10 numbers
rnsLaplace(n = 10, location = c(0.23, 0.24), scale = c(1, 2))
```

---

 Neutrosophic Negative Binomial

*Neutrosophic Negative Binomial Distribution*


---

**Description**

Density, distribution function, quantile function and random generation for the neutrosophic Negative Binomial distribution with parameters  $\text{size} = r_N$  and  $\text{prob} = p_N$ .

**Usage**

`dnsNegBinom(x, size, prob)`

`pnsNegBinom(q, size, prob, lower.tail = TRUE)`

`qnsNegBinom(p, size, prob)`

`rnsNegBinom(n, size, prob)`

**Arguments**

<code>x</code>	a vector or matrix of observations for which the pdf needs to be computed.
<code>size</code>	number of trials (zero or more), which must be a positive interval.
<code>prob</code>	probability of success on each trial, $0 < \text{prob} < 1$ .
<code>q</code>	a vector or matrix of quantiles for which the cdf needs to be computed.
<code>lower.tail</code>	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
<code>p</code>	a vector or matrix of probabilities for which the quantile needs to be computed.
<code>n</code>	number of random values to be generated.

**Details**

The neutrosophic negative binomial distribution with parameters  $r_N$  and  $p_N$  has the density

$$\binom{r_N + x - 1}{x} p_N^{r_N} (1 - p_N)^x$$

for  $r_N \in \{1, 2, \dots\}$  and  $p_N \in (p_L, p_U)$  which must be  $0 < p_N < 1$  and  $x \in \{0, 1, 2, \dots\}$ .

**Value**

`dnsNegBinom` gives the probability mass function

`pnsNegBinom` gives the distribution function

`qnsNegBinom` gives the quantile function

`rnsNegBinom` generates random variables from the Negative Binomial Distribution.

## References

Granados, C. (2022). Some discrete neutrosophic distributions with neutrosophic parameters based on neutrosophic random variables. *Hacetatepe Journal of Mathematics and Statistics*, 51(5), 1442-1457.

## Examples

```
dnsNegBinom(x = 1, size = 2, prob = c(0.5, 0.6))
pnsNegBinom(q = 1, size = 2, prob = c(0.5, 0.6))
qnsNegBinom(p = c(0.25, 0.5, 0.75), size = 2, prob = c(0.5, 0.6))
rnsNegBinom(n = 10, size = 2, prob = c(0.6, 0.6))
```

---

Neutrosophic Normal    *Neutrosophic Normal Distribution*

---

## Description

Density, distribution function, quantile function and random generation for the neutrosophic generalized exponential distribution with parameters mean =  $\mu_N$  and standard deviation sd =  $\sigma_N$ .

## Usage

```
dnsNorm(x, mean, sd)
pnsNorm(q, mean, sd, lower.tail = TRUE)
qnsNorm(p, mean, sd)
rnsNorm(n, mean, sd)
```

## Arguments

x	a vector or matrix of observations for which the pdf needs to be computed.
mean	the mean, which must be an interval.
sd	the standard deviations that must be positive.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

**Details**

The neutrosophic normal distribution with parameters mean  $\mu_N$  and standard deviation  $\sigma_N$  has density function

$$f_N(x) = \frac{1}{\sigma_N \sqrt{2\pi}} \exp\left\{-\frac{(X - \mu_N)^2}{2\sigma_N^2}\right\}$$

} for  $\mu_N \in (\mu_L, \mu_U)$ , the mean which must be an interval, and  $\sigma_N \in (\sigma_L, \sigma_U)$ , the standard deviation which must also be a positive interval, and  $-\infty < x < \infty$ .

**Value**

dnsNorm gives the density function

pnsNorm gives the distribution function

qnsNorm gives the quantile function

rnsNorm generates random variables from the neutrosophic normal distribution.

**References**

Patro, S. and Smarandache, F. (2016). The Neutrosophic Statistical Distribution, More Problems, More Solutions. Infinite Study.

**Examples**

```
data(balls)
dnsNorm(x = balls, mean = c(72.14087, 72.94087), sd = c(37.44544, 37.29067))

pnsNorm(q = 5, mean = c(72.14087, 72.94087), sd = c(37.44544, 37.29067))

# Calculate quantiles
qnsNorm(p = c(0.25, 0.5, 0.75), mean = c(9.1196, 9.2453), sd = c(10.1397, 10.4577))

# Simulate 10 values
rnsNorm(n = 10, mean = c(4.141, 4.180), sd = c(0.513, 0.521))
```

---

Neutrosophic Poisson    *Neutrosophic Poisson Distribution*

---

**Description**

Density, distribution function, quantile function and random generation for the neutrosophic Poisson distribution with parameter  $\lambda_N$ .

**Usage**

```
dnsPois(x, lambda)

pnsPois(q, lambda, lower.tail = TRUE)

qnsPois(p, lambda)

rnsPois(n, lambda)
```

**Arguments**

x	a vector or matrix of observations for which the pdf needs to be computed.
lambda	the mean, which must be a positive interval.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

**Details**

The neutrosophic Poisson distribution with parameter  $\lambda_N$  has the density

$$f_N(x) = \exp\{-\lambda_N\} \frac{(\lambda_N)^x}{x!}$$

for  $\lambda_N \in (\lambda_L, \lambda_U)$  which must be a positive interval and  $x \in \{0, 1, 2, \dots\}$ .

**Value**

dnsPois gives the probability mass function  
 pnsPois gives the distribution function  
 qnsPois gives the quantile function  
 rnsPois generates random variables from the neutrosophic Poisson Distribution.

**References**

Alhabib, R., Ranna, M. M., Farah, H., Salama, A. A. (2018). Some neutrosophic probability distributions. *Neutrosophic Sets and Systems*, 22, 30-38.

**Examples**

```
# In a company, Phone employee receives phone calls, the calls arrive with
# rate of [1 , 3] calls per minute, we will calculate
# the probability that the employee will not receive any call within a minute
dnsPois(x = 0, lambda = c(1, 3))

# the probability that employee would not receive any call within 5 minutes
dnsPois(x = 0, lambda = c(5, 15))
```

```

# the probability that the employee will receive at least one call within a minute
pnsPois(q = 1, lambda = c(1, 3), lower.tail = FALSE)
# the probability that the employee will receive at most three calls within 5 minutes
pnsPois(q = 3, lambda = c(5, 15), lower.tail = TRUE)
# Calcaute the quantiles
qnsPois(p = c(0.25, 0.5, 0.75), lambda = c(1, 3))
# Simulate 10 values
rnsPois(n = 10, lambda = 1)

```

---

### Neutrosophic Rayleigh *Neutrosophic Rayleigh Distribution*

---

#### Description

Density, distribution function, quantile function and random generation for the neutrosophic Rayleigh distribution with parameter  $\theta_N$ .

#### Usage

```

dnsRayleigh(x, theta)

pnsRayleigh(q, theta, lower.tail = TRUE)

qnsRayleigh(p, theta)

rnsRayleigh(n, theta)

```

#### Arguments

x	a vector or matrix of observations for which the pdf needs to be computed.
theta	the shape parameter, which must be a positive interval.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

#### Details

The neutrosophic Rayleigh distribution with parameter  $\theta_N$  has the density

$$f_N(x) = \frac{x}{\theta_N^2} \exp\left\{-\frac{1}{2} \left(\frac{x}{\theta_N}\right)^2\right\}$$

for  $\theta_N \in (\theta_L, \theta_U)$ , which must be a positive interval and  $x \geq 0$ .

**Value**

`dnsRayleigh` gives the density function

`pnsRayleigh` gives the distribution function

`qnsRayleigh` gives the quantile function

`rnsRayleigh` generates random variables from the Neutrosophic Rayleigh Distribution.

**References**

Khan, Z., Gulistan, M., Kausar, N. and Park, C. (2021). Neutrosophic Rayleigh Model With Some Basic Characteristics and Engineering Applications, in *IEEE Access*, 9, 71277-71283.

**Examples**

```
data(remission)
dnsRayleigh(x = remission, theta = c(9.6432, 9.8702))

pnsRayleigh(q = 20, theta = c(9.6432, 9.8702))

# Calculate quantiles
qnsRayleigh(p = c(0.25, 0.5, 0.75), theta = c(9.6432, 9.8702))

# Simulate 10 values
rnsRayleigh(n = 10, theta = c(9.6432, 9.8702))
```

---

Neutrosophic Uniform    *Neutrosophic Uniform Distribution*

---

**Description**

Density, distribution function, quantile function and random generation for the neutrosophic Uniform distribution of a continuous variable  $X$  with parameters  $a_N$  and  $b_N$ .

**Usage**

`dnsUnif(x, min, max)`

`pnsUnif(q, min, max, lower.tail = TRUE)`

`qnsUnif(p, min, max)`

`rnsUnif(n, min, max)`

**Arguments**

x	a vector or matrix of observations for which the pdf needs to be computed.
min	lower limits of the distribution. Must be finite.
max	upper limits of the distribution. Must be finite.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

**Details**

The neutrosophic Uniform distribution with parameters  $a_N$  and  $b_N$  has the density

$$f_N(x) = \frac{1}{b_N - a_N}$$

for  $a_N \in (a_L, a_U)$  lower parameter interval,  $b_N \in (b_L, b_U)$ , upper parameter interval.

**Value**

dnsUnif gives the density function

pnsUnif gives the distribution function

qnsUnif gives the quantile function

rnsUnif generates random variables from the neutrosophic Uniform Distribution.

**References**

Alhabib, R., Ranna, M. M., Farah, H., & Salama, A. A. (2018). Some neutrosophic probability distributions, *Neutrosophic Sets and Systems*, 22, 30-38.

**Examples**

```
dnsUnif(x = 1, min = c(0, 5), max = c(15, 20))
dnsUnif(x = c(6, 10), min = c(0, 5), max = c(15, 20))
```

```
punif(q = 1, min = c(0, 5), max = c(15, 20))
punif(q = c(6, 10), min = c(0, 5), max = c(15, 20))
```

```
qnsUnif(p = c(0.25, 0.5, 0.75), min = c(0, 5), max = c(15, 20))
```

```
rnsUnif(n = 10, min = c(0, 5), max = c(15, 20))
```

---

 Neutrosophic Weibull *Neutrosophic Weibull Distribution*


---

**Description**

Density, distribution function, quantile function and random generation for the neutrosophic Weibull distribution with scale parameter  $\alpha_N$  and shape parameter  $\beta_N$ .

**Usage**

```
dnsWeibull(x, shape, scale)
pnsWeibull(q, shape, scale, lower.tail = TRUE)
qnsWeibull(p, shape, scale)
rnsWeibull(n, shape, scale)
```

**Arguments**

x	a vector or matrix of observations for which the pdf needs to be computed.
shape	shape parameter, which must be a positive interval.
scale	scale parameter, which must be a positive interval.
q	a vector or matrix of quantiles for which the cdf needs to be computed.
lower.tail	logical; if TRUE (default), probabilities are $P(X \leq x)$ ; otherwise, $P(X > x)$ .
p	a vector or matrix of probabilities for which the quantile needs to be computed.
n	number of random values to be generated.

**Details**

The neutrosophic Rayleigh distribution with parameters  $\alpha_N$  and  $\beta_N$  has the density

$$f_N(x) = \frac{\beta_N}{\alpha_N^{\beta_N}} x^{\beta_N-1} \exp\{- (x/\alpha_N)^{\beta_N}\}$$

for  $\beta_N \in (\beta_L, \beta_U)$  the shape parameter must be a positive interval,  $\alpha_N \in (\alpha_L, \alpha_U)$ , the scale parameter which be a positive interval, and  $x > 0$ .

**Value**

dnsWeibull gives the density function  
 pnsWeibull gives the distribution function  
 qnsWeibull gives the quantile function  
 rnsWeibull generates random variables from the neutrosophic Weibull dDistribution.

## References

Alhasan, K. F. H. and Smarandache, F. (2019). Neutrosophic Weibull distribution and Neutrosophic Family Weibull Distribution, *Neutrosophic Sets and Systems*, 28, 191-199.

## Examples

```
data(remission)
dnsWeibull(x = remission, shape = c(1.0519, 1.0553), scale = c(9.3370, 9.4544))

pnsWeibull(q = 20, shape = c(1.0519, 1.0553), scale = c(9.3370, 9.4544))

# Calculate quantiles
qnsWeibull(p = c(0.25, 0.5, 0.75), shape = c(1.0519, 1.0553), scale = c(9.3370, 9.4544))

# Simulate 10 numbers
rnsWeibull(n = 10, shape = c(1.0519, 1.0553), scale = c(9.3370, 9.4544))
```

---

remission

*Remission data*

---

## Description

It is related to remission time in months of 128 cancer patients.

## Format

A data.frame with 128 observations of remission time in months of cancer patients.

## Source

Lee, E.T. and Wang, J. (2003), *Statistical Methods for Survival Data Analysis*. Vol. 476, John Wiley & Sons, Hoboken, NJ, USA.

Rao, G. S., Norouzirad, M., and Mazarei . D. (2023). Neutrosophic Generalized Exponential Distribution with Application. *Neutrosophic Sets and Systems*, 55, 471-485.

## Examples

```
data("remission")
remission
```

# Index

- balls, [2](#)
- beta, [3](#)
  
- dnsBeta (Neutrosophic Beta), [3](#)
- dnsBinom (Neutrosophic Binomial), [4](#)
- dnsDiscUnif (Neutrosophic Discrete Uniform), [6](#)
- dnsExp (Neutrosophic Exponential), [7](#)
- dnsGamma (Neutrosophic Gamma), [9](#)
- dnsGenExp (Neutrosophic Generalized Exponential), [10](#)
- dnsGenPareto (Neutrosophic Generalized Pareto), [11](#)
- dnsGenRayleigh (Neutrosophic Generalized Rayleigh), [13](#)
- dnsGeom (Neutrosophic Geometric), [14](#)
- dnsKumaraswamy (Neutrosophic Kumaraswamy), [16](#)
- dnsLaplace (Neutrosophic Laplace), [17](#)
- dnsNegBinom (Neutrosophic Negative Binomial), [19](#)
- dnsNorm (Neutrosophic Normal), [20](#)
- dnsPois (Neutrosophic Poisson), [21](#)
- dnsRayleigh (Neutrosophic Rayleigh), [23](#)
- dnsUnif (Neutrosophic Uniform), [24](#)
- dnsWeibull (Neutrosophic Weibull), [26](#)
  
- gamma, [9](#)
  
- Neutrosophic Beta, [3](#)
- Neutrosophic Binomial, [4](#)
- Neutrosophic Discrete Uniform, [6](#)
- Neutrosophic Exponential, [7](#)
- Neutrosophic Gamma, [9](#)
- Neutrosophic Generalized Exponential, [10](#)
- Neutrosophic Generalized Pareto, [11](#)
- Neutrosophic Generalized Rayleigh, [13](#)
- Neutrosophic Geometric, [14](#)
- Neutrosophic Kumaraswamy, [16](#)
- Neutrosophic Laplace, [17](#)
- Neutrosophic Negative Binomial, [19](#)
- Neutrosophic Normal, [20](#)
- Neutrosophic Poisson, [21](#)
- Neutrosophic Rayleigh, [23](#)
- Neutrosophic Uniform, [24](#)
- Neutrosophic Weibull, [26](#)
  
- pnsBeta (Neutrosophic Beta), [3](#)
- pnsBinom (Neutrosophic Binomial), [4](#)
- pnsDiscUnif (Neutrosophic Discrete Uniform), [6](#)
- pnsExp (Neutrosophic Exponential), [7](#)
- pnsGamma (Neutrosophic Gamma), [9](#)
- pnsGenExp (Neutrosophic Generalized Exponential), [10](#)
- pnsGenPareto (Neutrosophic Generalized Pareto), [11](#)
- pnsGenRayleigh (Neutrosophic Generalized Rayleigh), [13](#)
- pnsGeom (Neutrosophic Geometric), [14](#)
- pnsKumaraswamy (Neutrosophic Kumaraswamy), [16](#)
- pnsLaplace (Neutrosophic Laplace), [17](#)
- pnsNegBinom (Neutrosophic Negative Binomial), [19](#)
- pnsNorm (Neutrosophic Normal), [20](#)
- pnsPois (Neutrosophic Poisson), [21](#)
- pnsRayleigh (Neutrosophic Rayleigh), [23](#)
- pnsUnif (Neutrosophic Uniform), [24](#)
- pnsWeibull (Neutrosophic Weibull), [26](#)
  
- qnsBeta (Neutrosophic Beta), [3](#)
- qnsBinom (Neutrosophic Binomial), [4](#)
- qnsDiscUnif (Neutrosophic Discrete Uniform), [6](#)
- qnsExp (Neutrosophic Exponential), [7](#)
- qnsGamma (Neutrosophic Gamma), [9](#)
- qnsGenExp (Neutrosophic Generalized Exponential), [10](#)

qnsGenPareto (Neutrosophic Generalized Pareto), 11  
qnsGenRayleigh (Neutrosophic Generalized Rayleigh), 13  
qnsGeom (Neutrosophic Geometric), 14  
qnsKumaraswamy (Neutrosophic Kumaraswamy), 16  
qnsLaplace (Neutrosophic Laplace), 17  
qnsNegBinom (Neutrosophic Negative Binomial), 19  
qnsNorm (Neutrosophic Normal), 20  
qnsPois (Neutrosophic Poisson), 21  
qnsRayleigh (Neutrosophic Rayleigh), 23  
qnsUnif (Neutrosophic Uniform), 24  
qnsWeibull (Neutrosophic Weibull), 26

remission, 27

rnsBeta (Neutrosophic Beta), 3  
rnsBinom (Neutrosophic Binomial), 4  
rnsDiscUnif (Neutrosophic Discrete Uniform), 6  
rnsExp (Neutrosophic Exponential), 7  
rnsGamma (Neutrosophic Gamma), 9  
rnsGenExp (Neutrosophic Generalized Exponential), 10  
rnsGenPareto (Neutrosophic Generalized Pareto), 11  
rnsGenRayleigh (Neutrosophic Generalized Rayleigh), 13  
rnsGeom (Neutrosophic Geometric), 14  
rnsKumaraswamy (Neutrosophic Kumaraswamy), 16  
rnsLaplace (Neutrosophic Laplace), 17  
rnsNegBinom (Neutrosophic Negative Binomial), 19  
rnsNorm (Neutrosophic Normal), 20  
rnsPois (Neutrosophic Poisson), 21  
rnsRayleigh (Neutrosophic Rayleigh), 23  
rnsUnif (Neutrosophic Uniform), 24  
rnsWeibull (Neutrosophic Weibull), 26